


Systemic treatment of head and neck cancer

Tamara Rordorf

25.8.2017

- Role of chemotherapy in curative setting
- Recurrent/metastatic HNSCC: first line therapy and beyond
- Immunotherapy in HNSCC

Therapeutic concepts



T. Rordorf: Kopf-Hals Tumore

- Role of chemotherapy in curative setting (multimodal treatment)
 - primary (concurrent) radiochemotherapy
 - postoperative (adjuvant) radio(chemo)therapy
 - neoadjuvant chemotherapy
- r/mHNSCC: first line therapy and beyond
- Immunotherapy in HNSCC

LASCCHN: concomittant RCT



B.H, male, 68 y

LA HNSCC of hypopharynx cT3c, N2-3,G3
comorbidities: coronary heart disease

RT total dose 69.6 Gy
concomitant cisplatin 40 mg/m² weekly, for 5 weeks


Very good parital remission, rest FDG activity

Neck-dissection 10/10
follow up: complete remission




Pignon et al, 2009: Meta-analysis of chemotherapy in head and neck cancer (MACH-NC): An update on 93 randomised trials and 17,346 patients

(a) Concomitant chemotherapy.



(b) Induction chemotherapy



(a) Hazard ratio of death.

Timing	No. Deaths / No. Entered LRT+CT	No. Entered LRT	O-E	Variance	Hazard Ratio	HR [95% CI]
Concomitant	3171/4824	3389/4791	-326.4	1587.7		0.81 [0.78;0.86]
Induction	1877/2740	1813/2571	-40.0	900.7		0.96 [0.90;1.02]
Adjuvant	631/1244	661/1323	17.9	317.4		1.06 [0.95;1.18]
Total	5679/8808	5863/8685	-348.5	2805.8		0.88 [0.85;0.92]

Test for heterogeneity: $\chi^2_{107} = 179.8$ $p < 0.0001$

Test for interaction: $\chi^2_2 = 26.60$ $p < 0.0001$

$I^2 = 41\%$

LRT+CT better | LRT better

LRT+CT effect: $p < 0.0001$

concurrent chemotherapy: decrease in the risk of death
(HR 0.81, 95%CI 0.78-0.86)

Absolute survival benefit: 6.5 %

No survival benefit

for concurrent chemotherapy in pts over 70



for adjuvant or neoadjuvant chemotherapy

LASCCHN: primary surgery followed by adjuvant radiochemotherapy

79 yo male patient, current smoker

SCC floor of the mouth,, pT4a (3.5 cm), pN2c (7/42), M0, G2-3, L1, p16 positive;
surgery:

Mandibula resection, parital ressection of the tongue, reconstruction with fibular
transplant 6/14



-> ad RT oder RCT?

LAHNSCC: Adjuvante RT vs RCT

EORTC 22931

RTOG 95-01

ORIGINAL ARTICLE

Postoperative Irradiation with or without Concomitant Chemotherapy for Locally Advanced Head and Neck Cancer

Jacques Bernier, M.D., Ph.D., Christian Dornge, M.D., Mahmut Ozsahin, M.D., Ph.D., Katarzyna Matuszewska, M.D., Jean-Louis Lefebvre, M.D., Richard H. Greiner, M.D., Jordi Giral, M.D., Philippe Maingon, M.D., Frédéric Rolland, M.D., Michel Bolla, M.D., Francesco Cognetti, M.D., Jean Bourhis, M.D., Anne Kirkpatrick, M.Sc., and Martine van Glabbe, Ir., M.Sc., for the European Organization for Research and Treatment of Cancer Trial 22931

ABSTRACT

BACKGROUND

We compared concomitant cisplatin and irradiation with radiotherapy alone as adjuvant treatment for stage III or IV head and neck cancer.

METHODS

After undergoing surgery with curative intent, 167 patients were randomly assigned to receive radiotherapy alone (66 Gy over a period of 6½ weeks) and 167 to receive the same radiotherapy regimen combined with 100 mg of cisplatin per square meter of body-surface area on days 1, 22, and 43 of the radiotherapy regimen.

RESULTS

After a median follow-up of 60 months, the rate of progression-free survival was significantly higher in the combined-therapy group than in the group given radiotherapy alone ($P=0.04$ by the log-rank test; hazard ratio for disease progression, 0.75; 95 percent confidence interval, 0.56 to 0.99), with 5-year Kaplan-Meier estimates of progression-free survival of 47 percent and 36 percent, respectively. The overall survival rate was also significantly higher in the combined-therapy group than in the radiotherapy

The NEW ENGLAND JOURNAL of MEDICINE

ESTABLISHED IN 1812

MAY 6, 2004

VOL. 350 NO. 19

Postoperative Concurrent Radiotherapy and Chemotherapy for High-Risk Squamous-Cell Carcinoma of the Head and Neck

Jay S. Cooper, M.D., Thomas F. Pajak, Ph.D., Arlene A. Forastiere, M.D., John Jacobs, M.D., Bruce H. Campbell, M.D., Scott B. Saxman, M.D., Julie A. Kish, M.D., Harold E. Kim, M.D., Anthony J. Cmelak, M.D., Marvin Rotman, M.D., Mitchell Machtay, M.D., John F. Ensley, M.D., K.S. Clifford Chao, M.D., Christopher J. Schultz, M.D., Nancy Lee, M.D., and Karen K. Fu, M.D., for the Radiation Therapy Oncology Group 9501/Intergroup

ABSTRACT

BACKGROUND

Despite the use of resection and postoperative radiotherapy, high-risk squamous-cell carcinoma of the head and neck frequently recurs in the original tumor bed. We tested the hypothesis that concurrent postoperative administration of cisplatin and radiotherapy would improve the rate of local and regional control.

METHODS

Between September 9, 1995, and April 28, 2000, 459 patients were enrolled. After undergoing total resection of all visible and palpable disease, 231 patients were randomly assigned to receive radiotherapy alone (60 to 66 Gy in 30 to 33 fractions over a period of 6 to 6.6 weeks) and 228 patients to receive the identical treatment plus concurrent cisplatin (100 mg per square meter of body-surface area intravenously on days 1, 22, and 43).

RESULTS


After a median follow-up of 45.9 months, the rate of local and regional control was significantly higher in the combined-therapy group than in the group given radiotherapy alone (hazard ratio for local or regional recurrence, 0.61; 95 percent confidence interval,

From New York University Medical Center, New York (J.S.C.); Radiation Therapy Oncology Group Headquarters, Philadelphia (T.F.P.); Johns Hopkins Oncology Center, Baltimore (A.A.F.); Wayne State University School of Medicine, Detroit (J.J., H.E.K., J.F.E.); Medical College of Wisconsin, Milwaukee (B.H.C., C.J.S.); National Cancer Institute, Bethesda, Md. (S.B.S.); H. Lee Moffitt Cancer Center, Tampa, Fla. (J.A.K.); State University of New York Health Center at Brooklyn, Brooklyn (M.R.); University of Pennsylvania Health System, Philadelphia (M.M.); Maimonides Institute of Radiology, St. Louis (K.S.C.C.); and the University of California, San Francisco, San Francisco (N.L., K.K.F.).

N Engl J Med 2004;350:1937-44.

LASCC: postoperative RT vs RCT

Combined analysis of EORTC and RTOG 95-01 trial: only patients with ECE and R1 benefit from RCT (vs RT)



G.E., 1937. histology report

B 2014.30481 Neck dissection rechts Level II: Muskel- und Gewebsanteile ohne Lymphknotenstrukturen.

B 2014.30482 Neck dissection-Anteil III rechts: Metastase des Plattenepithelkarzinoms (0,4cm, ohne extrakapsuläres Wachstum) in einem Lymphknoten (1/1).

B 2014.30483 Neck dissection-Anteil IV rechts: Metastasen des Plattenepithelkarzinoms in zwei von zehn Lymphknoten (1,6 und 0,8cm, jeweils mit extrakapsulärem Wachstum) (2/10).

B 2014.30484 Neck dissection-Anteil V rechts: Sieben tumorfreie Lymphknoten (0/7).

B 2014.30485 Neck dissection Ib links: Zwei tumorfreie Lymphknoten (0/2).

B 2014.30486 Neck dissection II links: Metastase des Plattenepithelkarzinoms (1,3 cm, mit extrakapsulärem Wachstum) in einem von zwei Lymphknoten (1/2).

LAHNSCC: adjuvante RT vs RCT

RTOG 95-01 (Cooper et al, NEJM 2004)

- > RCT besser: 2 LK oder mehr; R1, oder extrakapsuläre Ausbreitung

EORTC trial (Bernier et al, NEJM 2004)

Combined Analysis


extranodal spread, R1 benefit from RCT vs RT

possible adverse risk factors:


vascular/lymphatic/perineural infiltration, pT3/4, oral/oropharynx with positive level 4 or 5 nodes: no benefit in joined analysis, benefit in EORTC trial (-> consider rCT)

EGFR signaling and inhibition

- epidermal growth factor receptor (EGFR) overexpressed in 90% of HNSCC; higher expression is associated with poor prognosis
- Activation: ligand binding on extracellular domain leads to activation of signalling pathways
- Blockade: monoclonal antibodies
- Cetuximab: chimeric IgG1 antibody against extracell. domain
- Improved OS when cetuximab added to standard of care
 - Phase 3 trial; combination with RT (curative)
 - Phase 3 trial; combination with chemotherapy (palliative disease)



LAHNSCC: RT +/- cetuximab («Bonner» Trial)



Cetuximab:

IgG1 chimerized antibody

Binds exclusively to EGFR and its heterodimers


Prevents repair and survival of tumor cells damaged by the effects of chemotherapy and radiotherapy

Potentiates apoptosis

Inhibits cell cycle progression

Decreases production of angiogenic factors

Inhibits invasion/metastasis



LAHNSCC: RT +/- cetuximab («Bonner» Trial)




Figure 2: Overall survival by treatment: 5-year update (median follow-up 60 months)




Figure 5: Overall survival by severity of rash in cetuximab-treated patients

	RT (n = 213)	RT + C (N=211)
Median survival,* mos	29.3	49
■ 95% confidence limits	21-38	36-58+
2 yrs, %	55	62
3 yrs, %	44	57
5 yrs, %	36.4	45.6
Log rank P value	.018	
HR (95% CI)	0.71 (0.54-0.95)	

LAHNSCC: Induction Chemotherapy

Primary chemotherapy followed by local curative treatment
(radio(chemo)therapy)

Rationale

HNSCC is chemotherapy-sensitive disease

-> up to 90% response rates in therapy-naive patients

radiation works better in smaller tumor volume

LAHNSCC: Induction Chemotherapy

O.C, 68 yo male, never smoker



oropharyngeal carcinoma; cT4cN2cN0, HPV positive, diagnosed in 1/10

difficulties swallowing, pain

Chemotherapy with docetaxel, cisplatin and 5FU (TCF)

pain and swallowing problems resolved after 1st cycle

curative RCT - > complete remission




Induction Chemotherapy

Selected Induction Taxane Trials

<u>Studies</u>	<u>Schema</u>	<u>Primary Objectives</u>
Tax 323 (Vermorken NEJM, 2007)		PFS, in stage III/IV M0 multisite "unresectable" SCC (N-358)
Tax 324 (Posner NEJM, 2007)		OS, in stage III/IV M0 multisite SCC (N-501)

TAX 324 Phase III : DPF vs PF followed by CT-RT Subset of Resectable Larynx/Hypopharynx SCC



But:

No arm included adequate chemoradiotherapy, other trials and meta-analysis showed no survival benefit compared to RCT

LAHNSCC: Unanswered questions

cisplatin schedule and dosis

is cetuximab equivalent to cisplatin?

role for cetuximab in adjuvant setting?

Is there a patient subroup that benefits from induction chemotherapy?

Platinum-based chemotherapy plus cetuximab in head and neck cancer.

Vermorken JB¹, Mesia R, Rivera F, Remenar E, Kaweck i A, Rottey S, Erfan J, Zabolotnyy D, Kienzer HR, Cupissol D, Pevrade E, Benasso M, Vynnychenko I, De Raucourt D, Bokemeyer C, Schueler A, Amellal N, Hitt R.




Table 2. Responses to Treatment and Survival.*

Variable	Cetuximab plus Platinum-Fluorouracil (N = 222)	Platinum-Fluorouracil Alone (N = 220)	Hazard Ratio or Odds Ratio (95% CI)	P Value
Survival — mo†				
Overall	10.1 (8.6–11.2)	7.4 (6.4–8.3)	Hazard ratio, 0.80 (0.64–0.99)	0.04‡
Progression-free	5.6 (5.0–6.0)	3.3 (2.9–4.3)	Hazard ratio, 0.54 (0.43–0.67)	<0.001‡
Best response to therapy — %				
Overall	36 (29–42)	20 (15–25)	Odds ratio, 2.33 (1.50–3.60)	<0.001§
Disease control¶	81 (75–86)	60.0 (53–67)	Odds ratio, 2.88 (1.87–4.44)	<0.001§
Time to treatment failure — mo†	4.8 (4.0–5.6)	3.0 (2.8–3.4)	Hazard ratio, 0.59 (0.48–0.73)	<0.001‡
Duration of response — mo	5.6 (4.7–6.0)	4.7 (3.6–5.9)	Hazard ratio, 0.76 (0.50–1.17)	0.21‡

- Role of chemotherapy in curative setting
- *r/mHNSCC: first line therapy and beyond*
- Immunotherapy in HNSCC: phase 3 data

Beyond the first line (until 2016):


Chemotherapy: taxanes, methotrexate, vinorelbine, cetuximab, monotherapy or combinations...

Selected chemotherapy trials

Study [ref]	Number of patients (# pts)	Treatment line/agent	RR	Disease stabilization/control rate	TPP	OS	Note
Grau ¹²	60	• Paclitaxel 80 mg/m ² weekly for 8 weeks; until progression for responders	43%	15%	8.2 months (responding patients only)		
Leon ¹³	151 total 43 chemotherapy 68 best supportive care (BSC) 40 radiotherapy	• Diverse	0	9.3%	N/A	107 days for chemotherapy group	
Cho ¹⁴	23	• Docetaxel 35 mg/m ² for 3 weeks of 4-week cycle, total 6 cycles	13%	34.7%	9 weeks	29 weeks	
Fayette ¹⁵	66	• Paclitaxel first line 48 pts • Paclitaxel second line 28 pts • Docetaxel 80 mg/m ² q3-4 w • Paclitaxel/carboplatin 175 mg/m ² /AUC 5q3w	30% 20% 16% 30%	36%	3.9 months	7.2 months	
Tahara ²²	74	• Paclitaxel 100 mg/m ² qw	29%	N/A	3.4	14.3	
Spece ²³	30	• Docetaxel 36 mg/m ² qw for 6 weeks	6.7%	33%	7.4 weeks	17.9 weeks	
Zenda ¹¹⁶	22	• Docetaxel 80 mg/m ² q3-4 w	10%	25%	1.7 months	4.6 months	
Nunico ¹¹⁷	18	• Docetaxel 80 mg/m ² q3w	11%	33%	19 weeks	26 weeks	
Vermorken ²⁵	103	• Eributix monotherapy followed by • Eributix-cis at progression	13%	46%	70 days	178 days	
Herbst ²⁴	51 with SD after cisplatin-based therapy 25 pts with progression under cisplatin 54 progressive within 90 days (amendment)	• Eributix/Cisplatin	18% 20% 6%		7.4 months 4.2 months 4.1 months	11.7 months 8.1 months 4.3 months	
Basile ²⁵	98 platinum-refractory	• Eributix/Cisplatin	10%	53%	85 days	183 days	
Machiels ²⁴	191	• Zalutumumab	6.3%	48%	9.9 weeks	6.7 months	
	95	• Best supportive care (BSC) (± Mtx)	1.1%	27%	8.4 weeks	5.2 months	


(RR: Response rate; TPP: Time to progression; OS: Overall survival; SD: Stable disease; AUC: Area under the concentration curve).

«targeted agents»



Afatinib versus methotrexate as second-line treatment in patients with recurrent or metastatic squamous-cell carcinoma of the head and neck progressing on or after platinum-based therapy (LUX-Head & Neck 1): an open-label, randomised phase 3 trial

Jean-Pascal H Machiels*, Robert I Haddad*, Jérôme Fayette*, Lisa F Licitra, Makoto Tahara, Jan B Vermorken, Paul M Clement, Thomas Gauler,




mPFS, months: 2.6 vs 1.7;
HR 0.80, P 0.03

Adverse events higher in afatinib arm:

Rash (10% in afatinib arm, vs 0)
Diarrhoea (9% vs 2%)
Stomatitis 6 % vs 8%

Buparlisib and paclitaxel in patients with platinum-pretreated recurrent or metastatic squamous cell carcinoma of the head and neck (BERIL-1): a randomised, double-blind, placebo-controlled phase 2 trial

Denis Soulières, Sandrine Favière, Ricard Mesia, Éva Remenár, Shau-Hsuan Li, Andrey Karpenko, Arunee Dechaphunkul, Sebastian Ochsenreither, Laura Anna Kiss, Jin-Ching Lin, Raj Nagarkar, László Tamás, Sung-Bae Kim, Jozsef Erfán, Anna Alyasova, Stefan Kasper, Carlo Barone, Sabine Turri, Arunava Chakravarty, Marie Chol, Paola Aimone, Samit Hirawat, Lisa Licitra



	Buparlisib and paclitaxel (n=79)	Placebo and paclitaxel (n=79)
Complete response	3 (4%)	1 (1%)
Partial response	28 (35%)	10 (13%)
Stable disease	26 (33%)	44 (56%)
Progressive disease	10 (13%)	19 (24%)
Unknown	11 (14%)	5 (6%)
Not assessed*	1 (1%)	0
Overall response†	31 (39%; 28.4–50.9)	11 (14%; 7.2–23.5)
Median duration of overall response (months)	4.5 (3.1–6.7)	7.1 (2.8–NE)
Disease control‡	57 (72%; 60.9–81.7)	55 (70%; 58.2–79.5)


Data are n (%), n (%; 95% CI), or median (95% CI). NE=not evaluable. * One patient did not have a baseline or post-baseline assessment. † Complete response and partial response. ‡ Complete response, partial response, or stable disease.

Table 2: Best overall response to therapy


mPFS: 4.6 months vs. 3.5 months; HR 0.65, p=0.01

- Chemotherapy in curative setting
- r/mHNSCC: first line therapy and beyond
- Immunotherapy in HNSCC

The immune system can eliminate cancer



Mutational load in carcinogen associated HNSCC



Alexandrov et al Nature 2013

Immunotherapy for HNSCC

Interleukin-2 given peritumoral, intranodal, intraarterial

Fishman Clin Cancer res. 2011; Rosenberg Ann surg.1989

Interleukin-12 intratumoral

Rapidis, JCO 2009

Interferon-gamma

conflicting data on tumor response

Richtsmeier WJ, Arch Otholaryngol 1990; Mahfoubi R, 1993, Head and neck

Monoclonal Antibodies

inhibit signalling pathway and initiate cell-mediated cytotoxicity

Immunotherapy for HNSCC

Conjugated Antibodies

mAb+ radioactive substance (Borjesson, Clin Cancer Res 2003)

Adoptive cell transfer

mostly in NPC (Smith et al 2012, Chia et al 2014)

patient's own ag-specific effector cells harvested; engineered and re-introduced to the patient (To, Arch Otolaryngol HN surgery 2000)

Anti-tumor vaccines

Dendritic-cell based (Bontkes Clin Immunol 2008, Gholamin BMC 2010)

peptide based (MAGE 3) (Voskens, 2012); whole tumor vaccines


Checkpoint inhibitors in HNSCC

Single substances:

- Nivolumab
- Pembrolizumab
- Darvolumab
- Avelumab

Combinations (clinical trials)

- Vaccines
- Immunotherapy/chemotherapy
- Two checkpoint inhibitors
- Radiation





The NEW ENGLAND
JOURNAL of MEDICINE

ORIGINAL ARTICLE

Nivolumab for Recurrent Squamous-Cell Carcinoma of the Head and Neck


R.L. Ferris, G. Blumenschein, Jr., J. Fayette, J. Guigay, A.D. Colevas, L. Licitra, K. Harrington, S. Kasper, E.E. Vokes, C. Even, F. Worden, N.F. Saba, L.C. Iglesias Docampo, R. Haddad, T. Rordorf, N. Kiyota, M. Tahara, M. Monga, M. Lynch, W.J. Geese, J. Kopit, J.W. Shaw, and M.L. Gillison

November 8 2016

Phase 3 CheckMate 141 Study Design

Nivolumab in R/M SCCHN After Platinum Therapy

Randomized, global, phase 3 trial of the efficacy and safety of nivolumab vs investigator's choice in patients with R/M SCCHN



Stratification factor


- Prior cetuximab treatment

^aTissue required for testing

DOR = duration of response; IV = intravenous; ORR = objective response rate; PFS = progression-free survival; Q2W = once every 2 weeks; R = randomized. Clinicaltrials.gov NCT02105636.


Overall Survival

Nivolumab in R/M SCCHN After Platinum Therapy



Progression-Free Survival

Nivolumab in R/M SCCHN After Platinum Therapy




Overall Survival by PD-L1 Expression

PD-L1 Expression $\geq 1\%$


Treatment Arm	Median OS, mo (95% CI)	HR (95% CI)
Nivolumab (n = 88)	8.7 (5.7-9.1)	0.55 (0.36-0.83)
Investigator's choice (n = 61)	4.6 (3.8-5.8)	

PD-L1 Expression $< 1\%$

Treatment Arm	Median OS, mo (95% CI)	HR (95% CI)
Nivolumab (n = 73)	5.7 (4.4-12.7)	0.89 (0.54-1.45)
Investigator's choice (n = 38)	5.8 (4.0-9.8)	



No. at Risk	0	3	6	9	12	15	18
Nivolumab	88	67	44	18	6	0	
Investigator's Choice	61	42	20	6	2	0	



No. at Risk	0	3	6	9	12	15	18
Nivolumab	73	52	33	17	8	3	0
Investigator's Choice	38	29	14	6	2	0	0

Ferris RL et al. *NEJM* 2016




Overall Survival by p16 (HPV) Status

p16-Positive


Treatment Arm	Median OS, mo (95% CI)	HR (95% CI)
Nivolumab (n = 63)	9.1 (7.2-10.0)	0.56 (0.32-0.99)
Investigator's choice (n = 29)	4.4 (3.0-9.8)	

p16-Negative

Treatment Arm	Median OS, mo (95% CI)	HR (95% CI)
Nivolumab (n = 50)	7.5 (3.0-NA)	0.73 (0.42-1.25)
Investigator's choice (n = 36)	5.8 (3.8-9.5)	




No. at Risk	63	49	35	18	10	3	0
Nivolumab	29	20	11	4	1	0	0
Investigator's Choice							



No. at Risk	50	32	25	12	6	1	0
Nivolumab	36	26	13	7	3	1	0
Investigator's Choice							

Ferris RL et al. *NEJM* 2016

EORTC QLQ-C30




Symptom Domains:

MID* ≥ 10 points

*10 points is conservative and smaller differences between groups are likely to be clinically relevant⁷

EORTC QLQ-H&N35




Minimally important difference ≥ 10 points

EORTC QLQ-C30 Functional Domains

CheckMate 141: Nivolumab vs IC in R/M SCCHN After Platinum Therapy

- Nivolumab-treated patients experienced stable PROs
- Investigator's choice–treated patients had statistically significant and clinically meaningful worsening in physical, role, and social functioning compared with nivolumab



Ongoing immunotherapy trials in HNSCC

Ongoing 2L Phase III trials in R/M HNSCC

Study	Population	Treatment Arms	N
KEYNOTE-040	R/M platinum refractory	<ul style="list-style-type: none"> Pembro 200 mg Q3W SoC (investigator's choice) 	600
EAGLE	R/M platinum refractory	<ul style="list-style-type: none"> MTX 40 mg/m² QW DOC 75 mg/m² Q3W CET 250 mg/m² QW Durvalumab monotherapy Durvalumab + Tremelimumab SoC 	720

Ongoing 1L Phase III trials in R/M HNSCC

Study	Population	Treatment Arms	N
CHECKMATE 651	Recurrent / metastatic	<ul style="list-style-type: none"> Nivo 3 mg/kg Q2W + Ipilimumab 1 mg/kg Q6W CET + 5-FU + plat 	490
KEYNOTE-048	Recurrent / metastatic	<ul style="list-style-type: none"> Pembro 200 mg Q3W Pembro + 5-FU + plat CET + 5-FU + plat 	780
KESTREL	Recurrent / metastatic	<ul style="list-style-type: none"> Durvalumab monotherapy Durvalumab + Tremelimumab SoC 	628

Summary

curative setting:

Cisplatin increases overall survival when given concurrently with radiotherapy
Cetuximab is an option for pat that can't receive cetuximab
in adjuvant setting, chemotherapy increases survival in pts with ENC and R1

metastatic disease:

- 1st line: Cetuximab, Platin and 5FU is the standard (and standard arm in the phase 3 immunotherapy-trials)
- After platinum-failure:
 - nivolumab is new standard (randomized phase 3 data showing survival benefit in patients who progressed after platinum-containing therapy)
- Multiple trials with immunotherapy in first and second lines; in combination with chemotherapy or other checkpoint inhibitors; oncolytic viruses

Thank you for your attention